

Solution of Fuzzy Volterra Integral Equations In A Bernstein Polynomial Basis

Maryam Mosleh

Department of Mathematics, Firoozkooh Branch, Islamic Azad University, Firoozkooh, Iran
Email: mosleh@iaufb.ac.ir

Mahmood Otadi

Department of Mathematics, Firoozkooh Branch, Islamic Azad University, Firoozkooh, Iran
Email: otadi@iaufb.ac.ir

Abstract— In this paper, we have used the parametric form of fuzzy number and convert a fuzzy Volterra integral equation to a system of integral equations in crisp case. We present a numerical method for solving fuzzy Volterra integral equations of the second kind. The proposed method is based on approximating unknown function with Bernstein's approximation. This method using simple computation with quite acceptable approximate solution. However, accuracy and efficiency are dependent on the size of the set of Bernstein polynomials. Furthermore we get an estimation of error bound for this method.

Index Terms— Fuzzy integral equations, System of Volterra integral equation, Bernstein polynomial

I. INTRODUCTION

The solutions of integral equations have a major role in the field of science and engineering. A physical even can be modelled by the differential equation, an integral equation. Since few of these equations cannot be solved explicitly, it is often necessary to resort to numerical techniques which are appropriate combinations of numerical integration and interpolation [1, 2]. There are several numerical methods for solving linear Volterra integral equation [3]. Kauthen in [4] used a collocation method to solve the Volterra- Fredholm integral equation numerically. Maleknejad and et al. in [5] obtained a numerical solution of Volterra integral equations by using Bernstein Polynomials.

The concept of fuzzy numbers and fuzzy arithmetic operations were first introduced by Zadeh [6], Dubois and Prade [7]. We refer the reader to [8] for more information on fuzzy numbers and fuzzy arithmetic. The topics of fuzzy integral equations (FIE) which growing interest for some time, in particular in relation to fuzzy control, have been rapidly developed in recent years. The fuzzy mapping function was introduced by Chang and Zadeh [9]. Later, Dubois and Prade [10] presented an elementary fuzzy calculus based on the extension principle also the concept of integration of fuzzy functions was first introduced by Dubois and Prade [10]. Babolian et al., Abbasbandy et al. in [11, 12] obtained a numerical solution of linear Fredholm fuzzy integral

equations of the second kind. Also, the fuzzy integral equations have been studied by several authors [13, 14, 15].

In this paper, we present a novel and very simple numerical method based upon Bernstein's approximation for solving Volterra fuzzy integral equations.

II. PRELIMINARIES

In this section the basic notations used in fuzzy calculus and Bernstein polynomials are introduced. We start by defining the fuzzy number.

Definition 1. [16] A fuzzy number is a fuzzy set $u : \mathbb{R}^1 \rightarrow I = [0,1]$ such that

- i. u is upper semi-continuous;
- ii. $u(x) = 0$ outside some interval $[a, d]$;
- iii. There are real numbers b and c , $a \leq b \leq c \leq d$,

for which

1. $u(x)$ is monotonically increasing on $[a, b]$,
2. $u(x)$ is monotonically decreasing on $[c, d]$,
3. $u(x) = 1, b \leq x \leq c$.

The set of all the fuzzy numbers (as given in definition 1) is denoted by E^1 .

An alternative definition which yields the same E^1 is given by Kaleva [17, 18].

Definition 2. A fuzzy number u is a pair (\underline{u}, \bar{u}) of functions $\underline{u}(r)$ and $\bar{u}(r)$, $0 \leq r \leq 1$, which satisfy the following requirements:

- i. $\underline{u}(r)$ is a bounded monotonically increasing, left continuous function on $(0,1]$ and right continuous at 0;
- ii. $\bar{u}(r)$ is a bounded monotonically decreasing, left continuous function on $(0,1]$ and right continuous at 0;
- iii. $\underline{u}(r) \leq \bar{u}(r), 0 \leq r \leq 1$.

A crisp number r is simply represented by $\underline{u}(\alpha) = \bar{u}(\alpha) = r, 0 \leq \alpha \leq 1$. The set of all the fuzzy numbers is denoted by E^1 .

For arbitrary $u = (\underline{u}(r), \bar{u}(r)), v = (\underline{v}(r), \bar{v}(r))$ and $k \in \mathbb{R}$ we define addition and multiplication by k as

$$\begin{aligned}(\underline{u} + \underline{v})(r) &= (\underline{u}(r) + \underline{v}(r)), \\ (\bar{u} + \bar{v})(r) &= (\bar{u}(r) + \bar{v}(r)), \\ k\underline{u}(r) &= k\underline{u}(r), k\bar{u}(r) = k\bar{u}(r), \text{ if } k \geq 0, \\ k\underline{u}(r) &= k\bar{u}(r), k\bar{u}(r) = k\underline{u}(r), \text{ if } k < 0.\end{aligned}$$

Remark 1. [12] Let $u = (\underline{u}(r), \bar{u}(r)), 0 \leq r \leq 1$ be a fuzzy number, we take

$$\begin{aligned}u^c(r) &= \frac{\underline{u}(r) + \bar{u}(r)}{2}, \\ u^d(r) &= \frac{\bar{u}(r) - \underline{u}(r)}{2}.\end{aligned}$$

It is clear that $u^d(r) \geq 0$, $\underline{u}(r) = u^c(r) - u^d(r)$ and $\bar{u}(r) = u^c(r) + u^d(r)$, also a fuzzy number $u \in E^1$ is said symmetric if $u^c(r)$ is independent of r for all $0 \leq r \leq 1$.

Remark 2. Let $u = (\underline{u}(r), \bar{u}(r)), v = (\underline{v}(r), \bar{v}(r))$ and also k, s are arbitrary real numbers. If $w = ku + sv$ then

$$\begin{aligned}w^c(r) &= ku^c(r) + sv^c(r), \\ w^d(r) &= |k| u^d(r) + |s| v^d(r).\end{aligned}$$

Definition 3. [19] For arbitrary fuzzy numbers u, v , we use the distance

$$D(u, v) = \sup_{0 \leq r \leq 1} \max\{|\bar{u}(r) - \bar{v}(r)|, |\underline{u}(r) - \underline{v}(r)|\}$$

and it is shown that (E^1, D) is a complete metric space [20].

The Bernstein's approximation, $B_n(f)$ to a real function $f : [0, 1] \rightarrow \mathbb{R}$ is the polynomial

$$B_n(f(x)) = \sum_{i=0}^n f\left(\frac{i}{n}\right) P_{n,i}(x), \quad (1)$$

where

$$P_{n,i}(x) = \binom{n}{i} x^i (1-x)^{n-i}, \quad i = 0, 1, \dots,$$

There are $n+1$ n th-degree polynomials. For convenience, we set $P_{n,i}(x) = 0$, if $i < 0$ or $i > n$. It can be readily shown that each of the Bernstein polynomials is positive.

Theorem 1. For all functions f in $C[0, 1]$, the sequence $B_n(f); n = 1, 2, 3, \dots$ converges uniformly to f , where B_n is defined by Eq. (1).

Proof. See [21].

This theorem follows that, for any $f \in C[0, 1]$ and for any $\varepsilon > 0$, there exists n such that the inequality $\|B_n(f) - f\| < \varepsilon$, holds.

We suppose $\|\cdot\|$ be the max norm on $[0, 1]$, then the error bound

$$\|B_n(f(x)) - f(x)\| \leq \frac{1}{2n} x(1-x) \|f''\|, \quad (2)$$

given in [22], shows that the rate of convergence is at least $\frac{1}{n}$ for $f \in C[0, 1]$. On the other hand, the asymptotic formula

$$\lim_{n \rightarrow \infty} n(B_n(f(x)) - f(x)) = \frac{1}{2} x(1-x) f''(x), \quad (3)$$

due to Voronovskaya [23] shows that for $x \in (0, 1)$

with $f' \neq 0$, the rate of convergence is precisely $\frac{1}{n}$.

III. FUZZY VOLTERRA INTEGRAL EQUATION

The Fuzzy Volterra integral equations of the second kind (FVIE-2) is [24]

$$F(x) = G(x) + \lambda \int_0^x k(x, t) F(t) dt, \quad 0 \leq x \leq 1. \quad (4)$$

where $\lambda > 0$, $k(x, t)$ is a kernel function and $G(x)$ is a fuzzy function. If $G(x)$ is a fuzzy function these equation may only possess fuzzy solution. Sufficient conditions for the existence of a unique solution to the fuzzy Volterra integral equation are given in [24].

Now, we introduce parametric form of a FVIE-2 with respect to Definition 2. Let $(\underline{G}(x; r), \bar{G}(x; r))$ and $(\underline{F}(x; r), \bar{F}(x; r)), 0 \leq r \leq 1$ are parametric form of $G(x)$ and $F(x)$, respectively then, parametric form of FVIE-2 is as follows:

$$\underline{F}(x; r) = \underline{G}(x; r) + \lambda \int_0^x k(x, t) \underline{F}(t; r) dt, \quad 0 \leq x \leq 1, \quad (5)$$

$$\bar{F}(x; r) = \bar{G}(x; r) + \lambda \int_0^x k(x, t) \bar{F}(t; r) dt. \quad (6)$$

Suppose $k(x, t)$ be continuous and for fix t , $k(x, t)$ changes its sign in finite points as x_j where $x_j \in [0, x]$. For example, let $k(x, t)$ be nonnegative over $[0, x_1]$ and negative over $[x_1, x]$, therefore from Eqs. (5) and (6), we have

$$\begin{aligned} \underline{F}(x;r) &= \underline{G}(x;r) + \lambda \int_0^x k(x,t) \underline{F}(t;r) dt \\ &+ \lambda \int_{x_1}^x k(x,t) \overline{F}(t;r) dt, 0 \leq x \leq 1, 0 \leq r \leq 1, \\ \overline{F}(x;r) &= \overline{G}(x;r) + \lambda \int_0^{x_1} k(x,t) \overline{F}(t;r) dt \\ &+ \lambda \int_{x_1}^x k(x,t) \underline{F}(t;r) dt, 0 \leq x \leq 1, 0 \leq r \leq 1. \end{aligned}$$

By referring to Remark 2 we have

$$\begin{aligned} F^c(x;r) &= G^c(x;r) + \lambda \int_0^x k(x,t) F^c(t;r) dt, \\ 0 \leq x \leq 1, \end{aligned} \quad (7)$$

$$F^d(x;r) = G^d(x;r) + \lambda \int_0^x |k(x,t)| F^d(t;r) dt, 0 \leq x \leq 1. \quad (8)$$

It is clear that we must solve two crisp Volterra integral equation of the second kind provided that each of Eqs. (7) and (8) have solution.

We consider the Volterra integral equations of the second kind given by,

$$\begin{aligned} F^c(x;r) &= G^c(x;r) + \lambda \int_0^x k(x,t) F^c(t;r) dt, \\ F^d(x;r) &= G^d(x;r) + \lambda \int_0^x |k(x,t)| F^d(t;r) dt, \\ 0 \leq x \leq 1, \end{aligned}$$

where $F^c(x;r)$ and $F^d(x;r)$ are the unknown crisp function to be determined, $k(x,t)$ is a continuous function on the square $[0,1]^2$ and integrable function, $G^c(x;r)$ and $G^d(x;r)$ being the known crisp functions.

To determine an approximate the unknown function of Eq. (4), we approximate with Bernstein's approximation

$$B_n(F(x)) = G(x) + \lambda \int_0^x k(x,t) B_n(F(t)) dt, \quad (9)$$

therefore, we approximate the unknown functions $F^c(x;r)$ and $F^d(x;r)$ by

$$B_n(F^c(x;r)) = \sum_{i=0}^n F^c\left(\frac{i}{n}; r\right) P_{n,i}(x), \quad (10)$$

and

$$B_n(F^d(x;r)) = \sum_{i=0}^n F^d\left(\frac{i}{n}; r\right) P_{n,i}(x), \quad (11)$$

where

$$P_{n,i}(x) = \binom{n}{i} x^i (1-x)^{n-i}, \quad i = 0, 1, \dots, n,$$

Let $(B_n(F(x;r)), \overline{B_n(F(x;r))})$, $0 \leq r \leq 1$ is a parametric form of $B_n(F(x))$, then we have:

$$\begin{aligned} \underline{B_n(F(x;r))} &= \sum_{i=0}^n F\left(\frac{i}{n}; r\right) P_{n,i}(x), 0 \leq r \leq 1, \\ \overline{B_n(F(x;r))} &= \sum_{i=0}^n F\left(\frac{i}{n}; r\right) P_{n,i}(x), 0 \leq r \leq 1 \end{aligned}$$

By referring to Remark 2, we have the following equations

$$\begin{aligned} \sum_{i=0}^n F^c\left(\frac{i}{n}; r\right) \binom{n}{i} (x^i (1-x)^{n-i} \\ - \lambda \int_0^x k(x,t) t^i (1-t)^{n-i} dt) = G^c(x;r), \\ 0 \leq r \leq 1, \end{aligned} \quad (12)$$

$$\begin{aligned} \sum_{i=0}^n F^d\left(\frac{i}{n}; r\right) \binom{n}{i} (x^i (1-x)^{n-i} \\ - \lambda \int_0^x |k(x,t)| t^i (1-t)^{n-i} dt), \end{aligned} \quad (13)$$

In order to find $F^c(\frac{i}{n}; r)$ and $F^d(\frac{i}{n}; r)$ for $i = 0, 1, \dots, n$, we now put $x = x_j$, $j = 0, 1, \dots, n$ in (12)

and (13), x_j 's being chosen as suitable distinct points in

$(0, 1]$, and x_0 is taken near 0 such that

$0 < x_0 < x_1 < \dots < x_n = 1$. Putting $x = x_j$ we obtain in short form two linear systems

$$A_1 X_1 = Y_1, \quad (14)$$

where

$$A_1 = \left[\binom{n}{i} (x_j^i (1-x_j)^{n-i}) \right]$$

$$- \lambda \int_0^{x_j} k(x_j, t) t^i (1-t)^{n-i} dt],$$

$$i, j = 0, 1, \dots, n,$$

$$X_1 = [F^c(\frac{i}{n}; r)]^t, \quad 0 \leq r \leq 1, i = 0, 1, \dots, n,$$

$$Y_1 = [G^c(x_j; r)]^t, \quad 0 \leq r \leq 1, j = 0, 1, \dots, n,$$

and also

$$A_2 X_2 = Y_2, \quad (15)$$

where

$$A_2 = \left[\binom{n}{i} (x_j^i (1-x_j)^{n-i} - \lambda \int_0^{x_j} |k(x_j, t)| t^i (1-t)^{n-i} dt) \right],$$

$$i, j = 0, 1, \dots, n,$$

$$X_1 = [F^d(\frac{i}{n}; r)]^t, \quad 0 \leq r \leq 1, i = 0, 1, \dots, n,$$

$$Y_1 = [G^d(x_j; r)]^t, \quad 0 \leq r \leq 1, j = 0, 1, \dots, n.$$

In general we cannot be able to carry out analytically the integrations, involved. We compute the integral that exist in A_1 's formula and A_2 's formula numerically. Now we

can show $F^c(\frac{i}{n}; r)$ and $F^d(\frac{i}{n}; r)$ by

$$F_n^c(\frac{i}{n}; r), i = 0, 1, \dots, n \text{ and } F_n^d(\frac{i}{n}; r), i = 0, 1, \dots, n,$$

respectively that are our solutions in nodes $x_j, j = 0, 1, \dots, n$ and by substituting them in Eqs. (10)

and (11) we can find $B_n(F_n^c(x_j; r))$ and $B_n(F_n^d(x_j; r)) j = 0, 1, \dots, n$ that are solution for integral equations (7) and (8).

We give error bound for this solution in the following theorem.

Theorem 2. Consider the crisp Volterra integral equations of the second kind (7) and (8). Assume that $k(x, t)$ is continuous on the square $[0, 1]^2$ and the solution of the equations belong to $(C^\alpha \cap L^2)([0, 1])$ for some $\alpha > 2$. If A_1 and A_2 invertible then

$$\sup_{x_i \in [0, 1]} D(F(x_i), B_n(F_n(x_i))) \leq$$

$$\sup_{r \in [0, 1]} \left[\frac{1}{8n} ((1+s) \|AA_1^{-1} \parallel F''^c(r) \| + \| F_n''^c(r) \|) \right]$$

$$\frac{1}{8n} ((1+s) \| A_2^{-1} \parallel F''^d(r) \| + \| F_n''^d(r) \|),$$

where $x_i = \frac{i}{n}, i = 0, 1, \dots, n, F(x)$ is exact solution of

FVIE-2 and $s = \sup_{x, t \in [0, 1]} |\lambda k(x, t)|$.

Proof. We have

$$\sup_{x_i \in [0, 1]} |F^\dagger(x_i; r) - B_n(F_n^\dagger(x_i; r))| =$$

$$\sup_{x_i \in [0, 1]} |F^\dagger(x_i; r) - F_n^\dagger(x_i; r) + F_n^\dagger(x_i; r) -$$

$$B_n(F_n^\dagger(x_i; r))| \leq$$

$$\sup_{x_i \in [0, 1]} |F^\dagger(x_i; r) - F_n^\dagger(x_i; r)| + \sup_{x_i \in [0, 1]}$$

$$|F_n^\dagger(x_i; r) - B_n(F_n^\dagger(x_i; r))|,$$

where \dagger means we have this equation for c and d together, independently. From relation (2) we have the following bound

$$\sup_{x \in [0, 1]} |F_n^\dagger(x; r) - B_n(F_n^\dagger(x; r))| \leq$$

$$\frac{1}{2n} x(1-x) \| F_n^\dagger \| \leq \frac{1}{8n} \| F_n^\dagger \|, (17)$$

then it is enough to find a bound for $\sup_{x_i \in [0, 1]} |F^\dagger(x_i; r) - F_n^\dagger(x_i; r)|$. For numerically

solving integral equations (7) and (8) by using Bernstein's approximation, because from Theorem 1 we

know that for any $F^\dagger \in C[0, 1]$ and for any $\varepsilon > 0$, there exists n such that the inequality

$\| B_n(F^\dagger) - F^\dagger \| < \varepsilon$, holds so we can write integral

equations (7) and (8) as

$$G^c(x; r) = B_n(F^c(x; r)) - \lambda \int_0^x k(x, t) B_n(F^c(t; r)) dt,$$

and

$$G^d(x; r) = B_n(F^d(x; r)) - \lambda \int_0^x |k(x, t)| B_n(F^d(t; r)) dt.$$

If we substitute $F_n^\dagger(x; r)$ instead of $F^\dagger(x; r)$ in above equation then the right-hand side of integral equation is exchanged by a new function that we denote it by $\hat{G}^\dagger(x; r)$. So we have,

$$\hat{G}^c(x; r) = B_n(F_n^c(x; r)) - \lambda \int_0^x k(x, t) B_n(F_n^c(t; r)) dt,$$

and

$$\hat{G}^d(x; r) = B_n(F_n^d(x; r)) - \lambda \int_0^x |k(x, t)| B_n(F_n^d(t; r)) dt,$$

Consequently we have

$$\sup_{x_i \in [0, 1]} |F^c(x_i; r) - F_n^c(x_i; r)| \leq \| A_1^{-1} \|$$

$$\max |G^c(x_i; r) - \hat{G}^c(x_i; r)|,$$

$$i = 0, 1, \dots, n,$$

(18)

$$\sup_{x_i \in [0, 1]} |F^c(x_i; r) - F_n^c(x_i; r)| \leq \| A_1^{-1} \|$$

$$\max |G^c(x_i; r) - \hat{G}^c(x_i; r)|,$$

$$i = 0, 1, \dots, n,$$

and

$$\sup_{x_i \in [0, 1]} |F^d(x_i; r) - F_n^d(x_i; r)| \leq \| A_2^{-1} \|$$

$$\max |G^d(x_i; r) - \hat{G}^d(x_i; r)|,$$

$$i = 0, 1, \dots, n,$$

(19)

where $x_i = \frac{i}{n}$, $i = 0, 1, \dots, n$. For finding a bound for $\max |G^\dagger(x_i; r) - \hat{G}^\dagger(x_i; r)|$,

we let

$$G^c(x; r) = F^c(x; r) - \lambda \int_0^x k(x, t) F^c(t; r) dt,$$

$$G^d(x; r) = F^d(x; r) - \lambda \int_0^x |k(x, t)| F^d(t; r) dt,$$

and

$$\hat{G}^c(x; r) = B_n(F^c(x; r)) - \lambda \int_0^x k(x, t) B_n(F^c(t; r)) dt,$$

$$\hat{G}^d(x; r) = B_n(F^d(x; r)) - \lambda \int_0^x |k(x, t)| B_n(F^d(t; r)) dt.$$

So that

$$G^c(x; r) - \hat{G}^c(x; r) = F^c(x; r) - B_n(F^c(x; r)) - \lambda \int_0^x k(x, t) (F^c(t; r) - B_n(F^c(t; r))) dt,$$

and

$$G^d(x; r) - \hat{G}^d(x; r) = F^d(x; r) - B_n(F^d(x; r)) - \lambda \int_0^x |k(x, t)| (F^d(t; r) - B_n(F^d(t; r))) dt,$$

if we let $\sup_{x, t \in [0, 1]} |\lambda k(x, t)| = s$, then we have

$$\begin{aligned} \sup |G^c(x; r) - \hat{G}^c(x; r)| &= \sup |F^c(x; r) - B_n(F^c(x; r)) - \\ &\lambda \int_0^x k(x, t) (F^c(t; r) - B_n(F^c(t; r))) dt| \leq \\ \sup |F^c(x; r) - B_n(F^c(x; r))| &+ \\ \sup |\lambda \int_0^x k(x, t) (F^c(t; r) - B_n(F^c(t; r))) dt| &\leq \\ B_n(F^c(x; r)) &+ \\ \sup |F^c(x; r) - B_n(F^c(x; r))| &+ \\ \sup |\lambda k(x, t) (F^c(t; r) - B_n(F^c(t; r)))| &\leq \\ \frac{1}{2n} x(1-x) \mathbf{P} F''^c \mathbf{P} + s \frac{1}{2n} t(1-t) & \\ \|F''^c\| \leq \frac{1}{8n} (1+s) \|F''^c\|, \end{aligned}$$

and

$$\sup |G^d(x; r) - \hat{G}^d(x; r)| \leq \frac{1}{8n} (1+s) \mathbf{P} F''^d \mathbf{P},$$

so by substituting this bound in the inequality (18) and (19) we have,

$$\sup_{x_i \in [0, 1]} |F^c(x_i; r) - F_n^c(x_i; r)| \leq \frac{1}{8n} (1+s) \|A_1^{-1}\| \|F''^c\|, \quad (20)$$

then from relations (16), (17), (20) and (21) we have

$$\sup_{x_i \in [0, 1]} |F^c(x_i; r) - B_n(F_n^c(x_i; r))| \leq \frac{1}{8n} ((1+s) \|A_1^{-1}\| \|F''^c\| + \|F_n''^c\|), \quad (22)$$

and

$$\begin{aligned} \sup_{x_i \in [0, 1]} |F^d(x_i; r) - F_n^d(x_i; r)| &\leq \\ \frac{1}{8n} (1+s) \|A_2^{-1}\| \|F''^d\|, & \\ \sup_{x_i \in [0, 1]} |F^d(x_i; r) - B_n(F_n^d(x_i; r))| &\leq \\ \frac{1}{8n} ((1+s) \|A_2^{-1}\| \|F''^d\| + \|F_n''^d\|), \end{aligned} \quad (23)$$

therefore by (22), (23) and Remark 1 we have

$$\begin{aligned} \sup_{x_i \in [0, 1]} |\overline{F(x_i; r)} - \overline{B_n(F_n(x_i; r))}| &\leq \\ \frac{1}{8n} ((1+s) \|A_1^{-1}\| \|F''^c\| + & \\ \|F_n''^c\|) + \frac{1}{8n} ((1+s) \|A_2^{-1}\| \|F''^d\| + & \\ \|F_n''^d\|), \end{aligned}$$

$$\begin{aligned} \sup_{x_i \in [0, 1]} |\overline{F(x_i; r)} - \overline{B_n(F_n(x_i; r))}| &\leq \\ \frac{1}{8n} ((1+s) \|A_1^{-1}\| \|F''^c\| + & \\ \|F_n''^c\|) + \frac{1}{8n} ((1+s) \|A_2^{-1}\| \|F''^d\| + & \\ \|F_n''^d\|), \end{aligned}$$

hence for all $r \in [0, 1]$

$$\begin{aligned} \max\{\sup_{x_i \in [0, 1]} |\overline{F(x_i; r)} - \overline{B_n(F_n(x_i; r))}|, & \\ |\overline{F(x_i; r)} - \overline{B_n(F_n(x_i; r))}|\} & \\ \leq \frac{1}{8n} ((1+s) \|A_1^{-1}\| \|F''^c\| + \|F_n''^c\|) & \\ + \frac{1}{8n} ((1+s) \|A_2^{-1}\| \|F''^d\| + \|F_n''^d\|), \end{aligned}$$

and then

and the proof is completed.

Theorem 3. (Minc [25]) The inverse of a nonnegative matrix A_2 is nonnegative if and only if A_2 is a generalized permutation matrix.

$$\sup_{x_i \in [0,1]} D(F(x_i), B_n(F_n(x_i))) \leq \sup_{r \in [0,1]} \left[\frac{1}{8n} ((1+s) \|A_1^{-1} \| \|F''^c(r)\| + \|F''^c(r)\|) + \frac{1}{8n} ((1+s) \|A_2^{-1} \| \|F''^d(r)\| + \|F''^d(r)\|) \right],$$

Theorem 4. Let $(G(x_j))$, $j = 0, 1, \dots, n$ is a fuzzy

$x_j^i(1-x_j)^{n-i} \geq \lambda \int_0^{x_j} |k(x_j, t)| t^i (1-t)^{n-i} dt$, $i, j = 0, 1, \dots, n$, arbitrary vector, and A_2 is a generalized permutation matrix, then Eq. (9) has a fuzzy Bernstein approximation.

$$x_j^i(1-x_j)^{n-i} \geq \lambda \int_0^{x_j} |k(x_j, t)| t^i (1-t)^{n-i} dt, i, j = 0, 1, \dots, n,$$

Proof. By we have $A_2 \geq 0$. By Theorem 3 and our hypotheses, proof is completed.

A. Comparison with Other Methods

In this subsection, the shortcomings of the existing methods [5, 12, 26] for solving fuzzy integral equations are pointed out.

Maleknejad and et al. in [5] obtained a numerical solution of Volterra integral equations of the form

$$F(x) = G(x) + \lambda \int_0^x k(x, t) F(t) dt, \quad 0 \leq x \leq 1$$

where $\lambda > 0$, $k(x, t)$ and $G(x)$ are real function, by using Bernstein Polynomials. But in this paper we consider more general form of fuzzy Volterra integral equation (4) where $\lambda > 0$, $k(x, t)$ is a kernel function and $G(x)$ is a fuzzy function.

Abbasbandy and et al. in [12] considered linear Fredholm fuzzy integral equations of the second kind of the form

$$F(x) = G(x) + \lambda \int_0^a k(x, t) F(t) dt, \quad 0 \leq x \leq 1$$

where $\lambda > 0$, $k(x, t)$ is a kernel function and $G(x)$ is a fuzzy function. In this paper, if we consider $x = a$ in equation (4) then this paper transfer to [12].

Abbasbandy and et al. in [26] used the homotopy analysis method (HAM) to obtain solution of fuzzy integro-differential equation

$$F'(x) + F(x) + \lambda \int_0^x k(x, t) F(t) dt = G(x), \quad 0 \leq x \leq 1.$$

But, in this paper we used Bernstein Polynomials to obtain solution of equation (4).

IV. NUMERICAL EXAMPLES

To illustrate the technique proposed in this paper, consider the following examples.

Example 4.1. We consider the fuzzy Volterra integral equation of the second kind given by,

$$F(x) = G(x) + \lambda \int_0^x F(t) dt, \quad 0 \leq x \leq 1,$$

where

$$\lambda = 1$$

$$G(x) = (\underline{G}(x; r), \overline{G}(x; r)) = (r, 2-r), \quad 0 \leq r \leq 1.$$

The exact solution in this case is given by

$$F(x) = (\underline{F}(x; r), \overline{F}(x; r)) = (re^x, (2-r)e^x), \quad 0 \leq r \leq 1.$$

We can see that

$$G^c(x; r) = 1, \quad G^d(x; r) = 1-r, \quad 0 \leq r \leq 1.$$

According to Eqs. (7) and (8) we have the following two crisp Volterra integral equations

$$F^c(x; r) = 1 + \int_0^x F^c(t; r) dt, \quad 0 \leq x \leq 1, \quad 0 \leq r \leq 1,$$

$$F^d(x; r) = 1-r + \int_0^x F^d(t; r) dt, \quad 0 \leq x \leq 1, \quad 0 \leq r \leq 1.$$

Now we approximate the unknown functions $F^c(x; r)$ and $F^d(x; r)$ by $B_n(F^c(x; r))$ and $B_n(F^d(x; r))$

TABLE I.
COMPUTED ERROR FOR EXAMPLES

n	Example 5.1	Example 5.2
	0.3513	0.5624
	0.0059	0.0015
	0.2844E-3	0.5848E-3

for $n = 1, 2, 3$.

We choose $x_0 = 10^{-10}$ and $x_j = \frac{j}{n}$ for $j = 1, \dots, n$, $n = 1, 2, 3$. For this example, we use

$r = 0, 0.1, \dots, 1$, where we calculate the error of the exact solution and obtained solution of fuzzy Volterra integral equation with Bernstein approximation. Table 1 show the convergence behavior for $n = 1, 2, 3$. The exact and obtained solution of fuzzy Volterra integral equation




Figure 1. Compares the exact solution and obtained solutions

in this example at $x = 0.5$ for $n = 1, 2, 3$, are shown in Figure 1.

Example 4.2. We consider the fuzzy Volterra integral equation of the second kind given by,

$$F(x) = G(x) + \lambda \int_0^x k(x, t) F(t) dt, \quad 0 \leq x \leq 1$$

where $\lambda = 1$, $k(x, t) = \sinh x$ and

$$G(x) = (\underline{G}(x; r), \overline{G}(x; r)) =$$

$$((\cosh x + 1 - \cosh^2 x)(r^2 + r),$$

$$(\cosh x + 1 - \cosh^2 x)(4 - r^3 - r)), 0 \leq r \leq 1.$$

The exact solution in this case is given by

$$F(x) = (\underline{F}(x; r), \overline{F}(x; r)) =$$

$$(\cosh x(r^2 + r), \cosh x(4 - r^3 - r)), 0 \leq r \leq 1.$$

We can see that

$$G^c(x; r) = \frac{(\cosh x + 1 - \cosh^2 x)(4 + r^2 - r^3)}{2},$$

$$G^d(x; r) = \frac{(\cosh x + 1 - \cosh^2 x)(4 - r^3 - r^2 - 2r)}{2}, \quad 0 \leq r \leq 1.$$

According to Eqs. (7) and (8) we have the following two crisp Volterra integral equations

$$F^c(x; r) = \frac{(\cosh x + 1 - \cosh^2 x)(4 + r^2 - r^3)}{2} + \int_0^x \sinh(x) F^c(t; r) dt, \quad 0 \leq x \leq 1, \quad 0 \leq r \leq 1,$$

$$F^d(x; r) = \frac{(\cosh x + 1 - \cosh^2 x)(4 - r^3 - r^2 - 2r)}{2} + \int_0^x \sinh(x) F^d(t; r) dt, \quad 0 \leq x \leq 1, \quad 0 \leq r \leq 1.$$

Now we approximate the unknown functions $F^c(x; r)$ and $F^d(x; r)$ by $B_n(F^c(x; r))$ and $B_n(F^d(x; r))$ for $n = 1, 2, 3$.

We choose $x_0 = 10^{-10}$ and $x_j = \frac{j}{n}$ for $j = 1, \dots, n$, $n = 1, 2, 3$. For this example, we use $r = 0, 0.1, \dots, 1$, where we calculate the error of the




Figure 2. Compares the exact solution and obtained solutions

exact solution and obtained solution of fuzzy Volterra integral equation with Bernstein approximation. Table 1 show the convergence behavior for $n = 1, 2, 3$. The exact and obtained solution of fuzzy Volterra integral equation in this example at $x = 0.5$ for $n = 1, 2, 3$, are shown in Figure 2.

V. CONCLUSIONS

Here a very simple and straight method, based on approximation of the fuzzy unknown function of an fuzzy Volterra integral equation on the Bernstein polynomial basis is used. Our achieve results in this paper, show that Bernstein's approximation method for solving fuzzy Volterra integral equations of second kind, is very effective and the answers are trusty and their accuracy are high and we can execute this method in a computer simply.

ACKNOWLEDGMENT

The authors are grateful to the anonymous referees for their valuable comments and suggestions to improve the presentation of this paper.

REFERENCES

- [1] C.T.H. Baker, "A perspective on the numerical treatment of Volterra equations," *J. Comput. Appl. Math.*, vol. 125, pp. 217-249, 2000.
- [2] P. Linz, "Analytical and numerical methods for Volterra equations," SIAM, Philadelphia, PA, 1985.
- [3] H.H. Sorkun and S. Yalcinbas, "Approximate solutions of linear Volterra integral equation systems with variable coefficients," *Applied Mathematical Modelling*, vol. 34, pp. 3451-3464, 2010.
- [4] J.P. Kauthen, "Continuous time collocation method for Volterra-Fredholm integral equations," *Numer. Math.*, vol. 56, pp. 409-424, 1989.
- [5] K. Maleknejad, E. Hashemizadeh and R. Ezzati, "A new approach to the numerical solution of Volterra integral equations by using Bernstein's approximation," *Commun Nonlinear Sci Numer Simulat*, vol. 16, pp. 647-655, 2011.
- [6] L.A. Zadeh, "The concept of a linguistic variable and its application to approximate reasoning," *Inform. Sci.*, vol. 8, pp. 199-249, 1975.
- [7] D. Dubois and H. Prade, "Operations on fuzzy numbers," *J. Systems Sci.*, vol. 9, pp. 613-626, 1978.
- [8] A. Kaufmann and M.M. Gupta, "Introduction Fuzzy Arithmetic, Van Nostrand Reinhold," New York, 1985.
- [9] S.S.L. Chang and L. Zadeh, "On fuzzy mapping and control," *IEEE Trans. System Man Cybernet*, vol. 2, pp. 30-34, 1972.
- [10] D. Dubois and H. Prade, "Towards fuzzdifferential calculus," *Fuzzy Sets Systems*, vol. 8, pp. 1-7, 1982.
- [11] E. Babolian, H.S. Goghary and S. Abbasbandy, "Numerical solution of linear Fredholm fuzzy integral equations of the second kind by Adomian method," *Applied Mathematics and Computation*, vol. 161, pp. 733-744, 2005.
- [12] S. Abbasbandy, E. Babolian and M. Alavi, "Numerical method for solving linear Fredholm fuzzy integral equations of the second kind," *Chaos Solitons & Fractals*, vol. 31, pp. 138-146, 2007.
- [13] R. Alikhani, F. Bahrami and A. Jabbari, "Existence of global solutions to nonlinear fuzzy Volterra integro-differential equations," *Nonlinear Analysis: Theory, Methods and Applications*, vol. 75, pp. 1810-1821, 2012.
- [14] A.M. Bica, "One-sided fuzzy numbers and applications to integral equations from epidemiology," *Fuzzy Sets and Systems*, vol. 219, pp. 27-48, 2013.
- [15] A.M. Bica and C. Popescu, "Numerical solutions of the nonlinear fuzzy Hammerstein-Volterra delay integral equations," *Information Sciences*, vol. 223, pp. 236-255, 2013.
- [16] G.J. Klir, U.S. Clair and B. Yuan, "Fuzzy set theory: foundations and applications," Prentice-Hall Inc., 1997.
- [17] O. Kaleva, "Fuzzy differential equations," *Fuzzy Sets Syst.*, vol. 24, pp. 301-317, 1987.
- [18] M. Ma, M. Friedman and A. Kandel, R.A. Devore, G.G. Lorentz
- [19] R. Goetschel and W. Vaxman, "Elementary calculus," *Fuzzy Sets Syst.*, vol. 18, pp. 31-43, 1986.
- [20] M.L. Puri and D. Ralescu, "Fuzzy random variables," *J. Math. Anal. Appl.*, vol. 114, pp. 409-422, 1986.
- [21] M.J.D. Powel, "Approximation theory and methods," Cambridge University Press, 1981.
- [22] R.A. Devore and G.G. Lorentz, "Constructive approximation," Berlin: Springer-Verlag, 1993.
- [23] E. Voronovskaya, "Determination de la forme asymptotique d'approximation des fonctions par les polynomes de M. Bernstein," *Doklady Akademii Nauk SSSR*, 1932.
- [24] J.Y. Park and H.K. Han, "Existence and uniqueness theorem for a solution of fuzzy Volterra integral equations," *Fuzzy Sets nad Systems*, vol. 105, pp. 481-488, 1999.
- [25] H. Minc, "Nonnegative matrices," Wiley, New York, 1988.
- [26] S. Abbasbandy and M.S. Hashemi, "A series solution of fuzzy integro-differential equations," *Journal of Fuzzy Set valued Analysis*, 1-17, 2012.

Maryam Mosleh borned in 1979 in Iran. She received the B.S., M.S., and Ph.D. degrees in applied mathematics from Islamic Azad University, Iran, in 2001, 2003 and 2009, respectively. She is currently an Associate Professor in the Department of Mathematics, Firoozkooh Branch, Islamic Azad University, Firoozkooh, Iran. She is actively pursuing research in fuzzy modeling, fuzzy neural network, fuzzy linear system, fuzzy differential equations and fuzzy integral equations. She has published numerous papers in this area.

Mahmood Otadi borned in 1978 in Iran. He received the B.S., M.S., and Ph.D. degrees in applied mathematics from Islamic Azad University, Iran, in 2001, 2003 and 2008, respectively. He is currently an Associate Professor in the Department of Mathematics, Firoozkooh Branch, Islamic Azad University, Firoozkooh, Iran. He is actively pursuing research in fuzzy modeling, fuzzy neural network, fuzzy linear system, fuzzy differential equations and fuzzy integral equations. He has published numerous papers in this area.

Call for Papers and Special Issues

Aims and Scope

JAIT is intended to reflect new directions of research and report latest advances. It is a platform for rapid dissemination of high quality research / application / work-in-progress articles on IT solutions for managing challenges and problems within the highlighted scope. JAIT encourages a multidisciplinary approach towards solving problems by harnessing the power of IT in the following areas:

- **Healthcare and Biomedicine** - advances in healthcare and biomedicine e.g. for fighting impending dangerous diseases - using IT to model transmission patterns and effective management of patients' records; expert systems to help diagnosis, etc.
- **Environmental Management** - climate change management, environmental impacts of events such as rapid urbanization and mass migration, air and water pollution (e.g. flow patterns of water or airborne pollutants), deforestation (e.g. processing and management of satellite imagery), depletion of natural resources, exploration of resources (e.g. using geographic information system analysis).
- **Popularization of Ubiquitous Computing** - foraging for computing / communication resources on the move (e.g. vehicular technology), smart / 'aware' environments, security and privacy in these contexts; human-centric computing; possible legal and social implications.
- **Commercial, Industrial and Governmental Applications** - how to use knowledge discovery to help improve productivity, resource management, day-to-day operations, decision support, deployment of human expertise, etc. Best practices in e-commerce, e-commerce, e-government, IT in construction/large project management, IT in agriculture (to improve crop yields and supply chain management), IT in business administration and enterprise computing, etc. with potential for cross-fertilization.
- **Social and Demographic Changes** - provide IT solutions that can help policy makers plan and manage issues such as rapid urbanization, mass internal migration (from rural to urban environments), graying populations, etc.
- **IT in Education and Entertainment** - complete end-to-end IT solutions for students of different abilities to learn better; best practices in e-learning; personalized tutoring systems. IT solutions for storage, indexing, retrieval and distribution of multimedia data for the film and music industry; virtual / augmented reality for entertainment purposes; restoration and management of old film/music archives.
- **Law and Order** - using IT to coordinate different law enforcement agencies' efforts so as to give them an edge over criminals and terrorists; effective and secure sharing of intelligence across national and international agencies; using IT to combat corrupt practices and commercial crimes such as frauds, rogue/unauthorized trading activities and accounting irregularities; traffic flow management and crowd control.

The main focus of the journal is on technical aspects (e.g. data mining, parallel computing, artificial intelligence, image processing (e.g. satellite imagery), video sequence analysis (e.g. surveillance video), predictive models, etc.), although a small element of social implications/issues could be allowed to put the technical aspects into perspective. In particular, we encourage a multidisciplinary / convergent approach based on the following broadly based branches of computer science for the application areas highlighted above:

Special Issue Guidelines

Special issues feature specifically aimed and targeted topics of interest contributed by authors responding to a particular Call for Papers or by invitation, edited by guest editor(s). We encourage you to submit proposals for creating special issues in areas that are of interest to the Journal. Preference will be given to proposals that cover some unique aspect of the technology and ones that include subjects that are timely and useful to the readers of the Journal. A Special Issue is typically made of 10 to 15 papers, with each paper 8 to 12 pages of length.

The following information should be included as part of the proposal:

- Proposed title for the Special Issue
- Description of the topic area to be focused upon and justification
- Review process for the selection and rejection of papers.
- Name, contact, position, affiliation, and biography of the Guest Editor(s)
- List of potential reviewers
- Potential authors to the issue
- Tentative time-table for the call for papers and reviews

If a proposal is accepted, the guest editor will be responsible for:

- Preparing the "Call for Papers" to be included on the Journal's Web site.
- Distribution of the Call for Papers broadly to various mailing lists and sites.
- Getting submissions, arranging review process, making decisions, and carrying out all correspondence with the authors. Authors should be informed the Instructions for Authors.
- Providing us the completed and approved final versions of the papers formatted in the Journal's style, together with all authors' contact information.
- Writing a one- or two-page introductory editorial to be published in the Special Issue.

Special Issue for a Conference/Workshop

A special issue for a Conference/Workshop is usually released in association with the committee members of the Conference/Workshop like general chairs and/or program chairs who are appointed as the Guest Editors of the Special Issue. Special Issue for a Conference/Workshop is typically made of 10 to 15 papers, with each paper 8 to 12 pages of length.

Guest Editors are involved in the following steps in guest-editing a Special Issue based on a Conference/Workshop:

- Selecting a Title for the Special Issue, e.g. "Special Issue: Selected Best Papers of XYZ Conference".
- Sending us a formal "Letter of Intent" for the Special Issue.
- Creating a "Call for Papers" for the Special Issue, posting it on the conference web site, and publicizing it to the conference attendees. Information about the Journal and Academy Publisher can be included in the Call for Papers.
- Establishing criteria for paper selection/rejections. The papers can be nominated based on multiple criteria, e.g. rank in review process plus the evaluation from the Session Chairs and the feedback from the Conference attendees.
- Selecting and inviting submissions, arranging review process, making decisions, and carrying out all correspondence with the authors. Authors should be informed the Author Instructions. Usually, the Proceedings manuscripts should be expanded and enhanced.
- Providing us the completed and approved final versions of the papers formatted in the Journal's style, together with all authors' contact information.
- Writing a one- or two-page introductory editorial to be published in the Special Issue.

More information is available on the web site at <http://www.academypublisher.com/jait/>.

